

Durham Research Online

Deposited in DRO:

16 August 2013

Version of attached file:

Accepted Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Carr, J.R. and Stokes, C.R. and Vieli, A. (2013) 'Recent progress in understanding marine-terminating Arctic outlet glacier response to climatic and oceanic forcing : twenty years of rapid change.', *Progress in physical geography*, 37 (4). pp. 435-466.

Further information on publisher's website:

<http://dx.doi.org/10.1177/0309133313483163>

Publisher's copyright statement:

The final definitive version of this article has been published in the Journal 'Progress in physical geography' 37/4, 2013
© SAGE Publications Ltd by SAGE Publications Ltd at the Progress in physical geography page:
<http://ppg.sagepub.com/> on SAGE Journals Online: <http://online.sagepub.com/>

Additional information:

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

Recent progress in understanding marine-terminating Arctic outlet glacier response to climatic and oceanic forcing: Twenty years of rapid change

Abstract

Until relatively recently, it was assumed that Arctic ice masses would respond to climatic/oceanic forcing over millennia, but observations made during the past two decades have radically altered this viewpoint and have demonstrated that marine-terminating outlet glaciers can undergo dramatic dynamic change at annual timescales. This paper reviews the substantial progress made in our understanding of the links between marine-terminating Arctic outlet glacier behaviour and the ocean-climate system during the past twenty years, when many ice masses have rapidly lost mass. Specifically, we assess three primary climatic/oceanic controls on outlet glacier dynamics, namely air temperature, ocean temperature and sea ice concentrations, and discuss key linkages between them. Despite recent progress, significant uncertainty remains over the response of marine-terminating outlet glaciers to these forcings, most notably (i), the spatial variation in the relative importance of each factor; (ii), the contribution of glacier-specific factors to glacier dynamics; and iii) the limitations in our ability to accurately model marine-terminating outlet glacier behaviour. Our present understanding precludes us from identifying patterns of outlet glacier response to forcing that are applicable across the Arctic and we

underscore the potential danger of extrapolating rates of mass loss from a small sample of study glaciers.

Keywords

Marine-terminating outlet glaciers, climate change, Arctic, cryosphere, Greenland Ice Sheet

I Introduction

Arctic warming is expected to far exceed the global average and is forecast to reach 4 to 7°C by 2100 (Meier et al., 2007; IPCC, 2007). Consequently, Arctic ice masses are expected to undergo rapid change during the 21st century and to contribute significantly to global sea level rise (e.g. Bamber et al., 2007). Indeed, estimates suggest that the Greenland Ice Sheet (GIS) contributed 0.46 mm a⁻¹ to sea level rise between 2000 and 2008 (van den Broeke et al., 2009). Assessing the potential response of Arctic ice masses to climate change is therefore crucial for the accurate prediction of near-future sea level rise (IPCC, 2007). For the purposes of this paper, we define ‘Arctic ice masses’ as the major glaciated archipelagos within the Arctic Circle, namely the Greenland Ice Sheet (GIS), Svalbard, Novaya Zemlya (NZ), Severnaya Zemlya (SZ), Franz Josef Land (FJL) and the Canadian Arctic (Figure 1). Alaska is also included as results from the region have contributed significantly to our knowledge of

marine-terminating outlet glacier dynamics. Here we define a marine-terminating outlet glacier as a channel of fast-moving ice that drains an ice cap or ice sheet and terminates in the ocean, at either a floating or grounded margin (Benn and Evans, 2010) (Figure 2).

Our understanding of Arctic ice mass behaviour has advanced dramatically during the last twenty years, particularly during the last decade. Previously, it was generally assumed that large Arctic ice masses would respond to climatic warming at millennial timescales, primarily through increased surface melting, and that changes in ice flow would occur only at centennial timescales or longer (Huybrechts et al., 1991; IPCC, 2001; Greve, 2000). However, studies published during the past two decades have dramatically altered this viewpoint (e.g. Joughin et al., 2010; Rignot et al., 2008; van den Broeke et al., 2009) and have shown that most Arctic ice masses have rapidly lost mass since the 1990s. Crucially, losses have been concentrated at the coastal margins, particularly on marine-terminating outlet glaciers (e.g. Meier et al., 2007; Thomas et al., 2009; Joughin et al., 2010). Indeed, recent studies have demonstrated that marine-terminating Arctic outlet glaciers can respond rapidly to climatic/oceanic forcing (e.g. Howat et al., 2007; Howat et al., 2008a; Joughin et al., 2008b; Howat et al., 2011; Joughin et al., 2010; Andersen et al., 2012; Kjær et al., 2012) and can significantly influence the mass budget of their parent ice masses over annual

to decadal timescales (e.g. Stearns and Hamilton, 2007; Pritchard et al., 2009; Rignot et al., 2008).

Results from the Antarctic, particularly Pine Island Glacier (Payne et al., 2004), have also highlighted the role of outlet glaciers and ice streams in enabling rapid coupling between forcing at the margins and the ice sheet interior and have raised concerns over the vulnerability of some regions to rapid mass loss (Joughin and Alley, 2011). Furthermore, iceberg-rafted debris from palaeo-ice sheets attest to major episodes of ice sheet collapse (e.g. Bond et al., 1992) and reconstructions of marine-based palaeo-ice sheets have highlighted the potential for rapid ice stream/outlet glacier retreat (Briner et al., 2009; e.g. Winsborrow et al., 2010). Theoretical considerations also suggest that glaciers resting on reverse bed slopes may potentially be unstable (Weertman, 1974; Thomas, 1979). Although this review focuses on the Arctic, these findings have demonstrated that marine-terminating outlet glaciers can respond rapidly to climatic/oceanic forcing and play a key role in regulating the mass balance of marine-based ice sheets. As a result, the factors controlling marine-terminating outlet glacier dynamics have emerged as a primary area of research.

Recent mass deficits have been attributed to both increased marine-terminating outlet glacier discharge and to a more negative surface mass balance (SMB), primarily resulting from increased surface melting relative to accumulation

(Rignot et al., 2011; van den Broeke et al., 2009; Rignot et al., 2008; Zwally et al., 2011). The relative contribution of each of these two components varies across the Arctic, but is presently approximately equal on the GIS (van den Broeke et al., 2009). A number of potential controls on marine-terminating outlet glacier behaviour have been identified (Figure 3), which we broadly classify as (i), glacier-specific factors, which relate to the glaciological, topographic and geological setting of the glacier; and (ii), climatic/oceanic forcing, including air and ocean temperatures, sea ice and precipitation. Important glacier-specific factors include subglacial topography and geology, fjord bathymetry and topography, sedimentation at the grounding line and glacier velocity, size, surface slope and catchment area (Figure 3) (Meier and Post, 1987; Alley, 1991; Joughin et al., 2008b). Theory suggests that changes in marine-terminating outlet glacier dynamics can occur independently of climatic/oceanic forcing (e.g. Alley, 1991; Meier and Post, 1987) and the importance of glacier-specific factors, particularly subglacial topography, has been highlighted by recent studies (Thomas et al., 2009; Joughin et al., 2010; Joughin et al., 2012). Despite their apparent significance, however, the influence of glacier-specific factors on Arctic marine-terminating glacier behaviour is poorly understood.

In contrast, concerns over anthropogenic climate change in the 1990s resulted in an increasing focus on climatic/oceanic forcing factors and recent work has

emphasised the widespread and synchronous nature of dynamic changes in many regions, particularly south-eastern Greenland (e.g. Howat et al., 2008a; Murray et al., 2010). Consequently, this paper focuses on the climatic/oceanic drivers of marine-terminating Arctic outlet glacier dynamics and discusses three primary controls: air temperatures, ocean temperatures and sea ice concentrations (Figure 3). It should be noted, however, that these forcing factors are not independent (Figure 3) and that interconnections between them may significantly influence outlet glacier behaviour, yet many of these relationships are poorly understood. We aim to: i), review and summarise recent developments relating to each of these climatic/oceanic forcing factors; ii), highlight key uncertainties surrounding marine-terminating Arctic outlet glacier response to climatic/oceanic forcing; and iii), recommend directions for future research.

II Arctic mass balance trends: 1990 to 2010

Rapid mass loss from Arctic masses has been documented since the early 1990s by numerous independent studies (Table 1) (e.g. Moholdt et al., 2010b; Krabill et al., 2004; Rignot and Kanagaratnam, 2006; Velicogna and Wahr, 2006; Gardner et al., 2011). Due to their remote location and considerable size, mass balance is usually determined indirectly using remotely sensed data and/or SMB modelling. Considerable advances have been made in these techniques

during the past twenty years, which have substantially improved our ability to quantify mass budgets and to assess the relative contribution of ice dynamics to mass loss (van den Broeke et al., 2009; Rignot and Kanagaratnam, 2006; Velicogna and Wahr, 2006; Krabill et al., 2004). At present, the primary techniques include Gravity Recovery and Climate Experiment (GRACE) data (e.g. Luthcke et al., 2006; Velicogna, 2009; Velicogna and Wahr, 2006; Khan et al., 2010; Mémin et al., 2011; Arendt et al., 2008; Wouters et al., 2008; Jacob et al., 2012; Bergmann et al., 2012), comparison of SMB with outlet glacier discharge (Rignot et al., 2008; Rignot and Kanagaratnam, 2006; van den Broeke et al., 2009; Rignot et al., 2011) and repeat laser or radar altimetry measurements (Krabill et al., 2004; Thomas et al., 2006; Abdalati et al., 2001; Thomas et al., 2009; Pritchard et al., 2009).

The negative mass balance of the GIS has received particular attention and has been estimated via a number of techniques and for a range of time periods. The most recent values from GRACE (Jacob et al., 2012) and from the comparison of SMB/outlet glacier discharge (Rignot et al., 2011) are presented in Table 1. An important new trend is the rapid mass loss from the Canadian Arctic between 2007 and 2009, which made the archipelago the primary cryospheric contributor to eustatic sea level rise outside of the Greenland and Antarctic ice sheets (Table 1) (Gardner et al., 2011). Furthermore, the area has been

highlighted as the largest potential contributor to ice loss and sea level rise of any glaciated region during the 21st century (Radić and Hock, 2011). Negative mass balance trends have also been documented in Svalbard (Nuth et al., 2010; Hagen et al., 2009; Moholdt et al., 2010b) and the Russian Arctic (Table 1) (Sharov et al., 2009; Kotlyakov et al., 2010). However, the mass balance of the Russian Arctic archipelagos have been comparatively poorly documented (Bassford et al., 2006). This represents a significant limitation to our understanding of the Arctic cryosphere and highlights the need for further research in the region, as NZ, SZ and FJL account for approximately 20% of the glaciated area of the Arctic, excluding the GIS (Dowdeswell et al., 1997).

1 Spatial trends in Arctic mass balance

Arctic mass balance trends have been spatially non-uniform, with many areas exhibiting slight growth at high elevations and rapid marginal thinning (e.g. Thomas et al., 2008; Hagen et al., 2009; Sharov et al., 2009; Thomas et al., 2006; Pritchard et al., 2009; Zwally et al., 2011; Sharov, 2010). Substantial thickening has been observed at high elevations on the GIS (Johannessen et al., 2005; Thomas et al., 2006; Zwally et al., 2005; Ettema et al., 2009); Austfonna ice cap, Svalbard (Moholdt et al., 2010a; Bamber et al., 2004; Raper et al., 2005; Moholdt et al., 2010b); the northern ice cap, NZ (Sharov et al., 2009); Tyndall and Windy ice domes in FJL; Schmidt and Vavilov ice caps in SZ (Sharov,

2010); and some Canadian Arctic ice caps (Mair et al., 2009; Abdalati et al., 2004). A number of potential explanations have been proposed for this interior thickening, including increased precipitation (Thomas et al., 2006; Zwally et al., 2005), possibly related to changes in sea ice extent (Mair et al., 2009; Bamber et al., 2004; Raper et al., 2005), long-term accumulation trends (Koerner, 2005; Moholdt et al., 2010a) and/or surge dynamics (Bevan et al., 2007). However, interior gains have been far outweighed by low-elevation thinning and marginal retreat (e.g. Zwally et al., 2011; van den Broeke et al., 2009), resulting in an overall negative mass balance in many regions (Table 1).

2 Dynamic contribution of marine-terminating outlet glaciers to mass loss

In addition to rapid marginal thinning, peak losses have occurred on marine-terminating outlet glaciers (Pritchard et al., 2009; Moon and Joughin, 2008; Sole et al., 2008). On many of these glaciers, thinning rates of 10s of m a^{-1} have far exceeded surface melt rates, suggesting that thinning is largely 'dynamic' (i.e. resulting from changes in ice flow, rather than increased surface melting) (e.g. Abdalati et al., 2001; Krabill et al., 2004; Burgess and Sharp, 2008; Thomas et al., 2009). The contribution of glacier dynamics to recent mass deficits has been further emphasised by rapid retreat rates, which have reached kilometres per year on the GIS (e.g. Howat et al., 2008a; Moon and Joughin, 2008; Joughin et al., 2008b; Joughin et al., 2010) and hundreds of metres per year elsewhere

(e.g. Sharov, 2005; Blaszczyk et al., 2009; Nuth et al., 2010; Burgess and Sharp, 2004). Furthermore, recent research has underscored the contribution of dynamic changes to decadal-scale losses, as initial perturbations at the glacier terminus may be rapidly transmitted to inland areas, producing widespread, substantial thinning (Zwally et al., 2011; Pritchard et al., 2009; Thomas et al., 2011; Howat et al., 2008b; Howat et al., 2005). This longer-term component of dynamic loss is an important emerging area of research and has the potential to be the primary component of the GIS contribution to 21st century sea level rise (Price et al., 2011; Vieli and Nick, 2011).

Although the dynamics of marine-terminating outlet glaciers are now recognised as a key component of Arctic ice mass loss, they have also been highlighted as a principle area of uncertainty (IPCC, 2007). Specifically, the primary climatic/oceanic controls and the mechanisms by which they induce a dynamic response are yet to be fully understood (Vieli and Nick, 2011; Sole et al., 2008; Howat et al., 2010). The following sections review the three main climatic/oceanic controls identified to date, namely surface air temperatures, ocean temperatures and sea ice concentrations, and discuss the primary linkages between these factors (Figure 3). All three forcing factors have undergone marked changes in recent years, which have been linked to both recent climatic warming (IPCC, 2007; ACIA, 2004) and to the onset of a

negative phase of the North Atlantic Oscillation (NAO) in the mid-1990s (Stern and Heide-Jørgensen, 2003; Gerdes et al., 2003; Hurrell et al., 2003; e.g. Holliday et al., 2008).

III Air temperature forcing

Arctic air temperatures have risen substantially since the mid-1990s (Hanna et al., 2008; IPCC, 2007; ACIA, 2004), although they are not unprecedented at decadal timescales (Chylek et al., 2006; Box et al., 2009). We present a new synthesis of air temperature data to investigate the spatial distribution of Arctic warming between 1990 and 2010 and to visualise this trend both in terms of magnitude and statistical significance (Figure 4). Linear trends were calculated from annual air temperature series, which were compiled from meteorological station data of varying temporal resolution (three-hourly to monthly). In order to account for missing values, three-hourly data were used only if: i), no more than two consecutive records were missing in a day and; ii), no more than three records in total were missing in a day. Daily data were only used if values were available for 22 or more days per month and monthly values were used only if data were available for all months of the year (Cappelen, 2011).

Results suggest that warming has been greatest at coastal stations surrounding Baffin Bay and the Davis Strait (Figure 4), which is consistent with dramatic

mass loss from the Canadian Arctic between 2004 and 2009 (Gardner et al., 2011). Significant warming has also occurred in the Kara Sea region, particularly on FJL (Figure 4), although data coverage is comparatively sparse. Warming from the mid-1990s has been linked to negative SMB on a number of Arctic ice masses, particularly the GIS (e.g. Bhattacharya et al., 2009; Hanna et al., 2008; Box et al., 2006; Mote, 2007; Ettema et al., 2009; Abdalati and Steffen, 2001). However, whilst warming directly affects SMB, a key recent development has been to consider the potential impact of meltwater on outlet glacier dynamics.

1 Air temperatures, meltwater production and ice velocities on temperate and polythermal glaciers

The relationship between air temperatures, meltwater supply and ice velocities has been well-documented on temperate glaciers (e.g. Fountain and Walder, 1998; Iken and Bindshadler, 1986; Willis, 1995), but had not been extensively considered on large Arctic ice masses until relatively recently. On temperate glaciers, surface meltwater is thought to access large portions of the glacier bed during the melt season, resulting in elevated basal water pressures, reduced basal drag and enhanced ice motion (e.g. Fountain and Walder, 1998; Nienow et al., 1998; Willis, 1995; Iken and Bindshadler, 1986; Kamb, 1987). As the melt season progresses, continued meltwater input promotes the development

of a more efficient subglacial drainage system, which lowers basal water pressures and reduces the sensitivity of glacier velocities to additional melt (Figure 5) (e.g. Nienow et al., 1998; Willis, 1995). Recent studies have demonstrated a similar relationship on polythermal glaciers in the Canadian Arctic (e.g. Copland et al., 2003; Bingham et al., 2008; Bingham et al., 2003; Boon and Sharp, 2003) and in Svalbard (Rippin et al., 2005; Vieli et al., 2004; Nuttall and Hodgkins, 2005). In particular, extensive investigations on John Evans Glacier (JEG), Ellesmere Island, Canada, showed that surface meltwater could rapidly access the bed through predominantly cold ice and cause substantial seasonal acceleration (Copland et al., 2003; Bingham et al., 2008; Bingham et al., 2003; Bingham et al., 2005).

2 Surface meltwater and ice velocities in the GIS ablation zone

Until a decade ago, it was largely assumed that penetration of surface meltwater to the bed of large Arctic ice masses would be minimal and that its effect on ice velocities would be limited, especially on the GIS (Hodgkins, 1997; Copland et al., 2003; Zwally et al., 2002). This viewpoint was radically altered by GPS measurements from Swiss Camp in the west Greenland ablation zone, which first demonstrated a close correspondence between surface meltwater inputs and ice velocities (Zwally et al., 2002). Here we define the ablation zone as areas that experience melt, with the exception of fast-flowing, marine

terminating outlet glaciers, which are discussed separately (Section II 3), due to their differing response to meltwater inputs. Results from Swiss Camp showed that velocities closely followed seasonal and interannual variations in surface meltwater production, as previously observed on temperate glaciers, and this was attributed to meltwater-enhanced basal sliding (Zwally et al., 2002). Most importantly, the study highlighted meltwater-enhanced basal lubrication as a potential mechanism for rapid, dynamic and widespread response of the GIS to atmospheric warming (Zwally et al., 2002).

The work of Zwally *et al.* (2002) was supported by subsequent results from the west Greenland ablation zone, which provided further evidence of rapid coupling between seasonal meltwater inputs and ice velocities (e.g. Catania and Neumann, 2010; Bartholomew et al., 2010; Bartholomew et al., 2011; Das et al., 2008; Joughin et al., 2008a; van de Wal et al., 2008). Studies also identified supraglacial lake drainage events as a potential mechanism for rapid transfer of meltwater to the bed (e.g. Krawczynski et al., 2009; Das et al., 2008). Large volumes of water released during drainage events may promote crevasse propagation through the full ice thickness by offsetting rapid refreezing and maintaining high water pressures at the crevasse tip (Krawczynski et al., 2009; van der Veen, 2007; Alley et al., 2005; van der Veen, 1998). Drainage events have immediately preceded velocity increases in the west Greenland ablation

zone (Das *et al.*, 2008; Box and Ski, 2007; McMillan *et al.*, 2007), on land-terminating west Greenland outlet glaciers (Sneed and Hamilton, 2007; Shepherd *et al.*, 2009) and on JEG (Copland *et al.*, 2003; Bingham *et al.*, 2003; Boon and Sharp, 2003), providing empirical support for their role in meltwater delivery to the bed.

The potential impact of surface meltwater inputs on the GIS was also explored using numerical modelling, which predicted far greater losses with enhanced basal sliding (Parizek and Alley, 2004; Huybrechts and de Wolde, 1999; van de Wal and Oerlemans, 1997). This occurred via a number of proposed feedback mechanisms, which are illustrated for an idealised section of the GIS (Figure 6). Specifically, feedbacks could develop between glacier acceleration, dynamic thinning and surface melting: increased basal sliding would promote dynamic thinning and bring a greater portion of the ice sheet into the ablation zone, thus exposing a greater area to melting and enhanced lubrication (Figure 6) (Parizek and Alley, 2004).

3 Surface meltwater and marine-terminating Arctic outlet glacier dynamics

The close coupling between surface meltwater and ice velocities observed in the GIS ablation zone led to increased consideration of the influence of meltwater on marine-terminating outlet glacier dynamics (e.g. Hall *et al.*, 2008; Krabill *et al.*, 2004). This was further motivated by the concurrence of the onset

of marine-terminating Arctic glacier retreat from the mid-1990s with atmospheric warming (e.g. Howat and Eddy, 2011; Dyurgerov and McCabe, 2006; Bevan et al., 2012a) and the coincidence of substantial changes in glacier dynamics with elevated air temperatures (e.g. Howat et al., 2008a; Moon and Joughin, 2008; Rignot and Kanagaratnam, 2006).

Recent results from marine-terminating Arctic outlet glaciers appear to support meltwater-enhanced basal lubrication as a mechanism for ice acceleration at sub-annual timescales: glacier velocities in the Uummannaq region of west Greenland (Howat et al., 2010) and on Duvebreen, Austfonna (Dunse et al., 2012) (Figure 1), closely corresponded to the seasonal melt cycle. Similarly, results from Petermann Glacier (Figures 1 & 2) (Nick et al., 2012) and Dagaard Jensen Gletscher (Figure 1) (Bevan et al., 2012b) suggest that seasonal velocities primarily reflect variations in surface meltwater availability and data from Helheim Glacier (HH) (Figure 1) indicate that surface meltwater can be transmitted to the bed within 12 to 36 hours (Andersen et al., 2010a).

Despite an apparent relationship at seasonal or shorter timescales, however, the influence of meltwater-enhanced basal lubrication on interannual marine-terminating outlet glacier behaviour remains equivocal (e.g. Bingham et al., 2003; Vieli et al., 2004; van de Wal et al., 2008; Seale et al., 2011; McFadden et al., 2011). Evidence from the GIS suggests that meltwater input to the bed may

have a limited impact on interannual velocity changes on fast-flowing marine-terminating outlet glaciers and that ice flow may be more responsive to conditions at the ice-ocean interface (Joughin et al., 2008a; Nick et al., 2009). A similar pattern has been observed on JEG (Bingham et al., 2003) and Hansbreen, Spitzbergen (Figure 1) (Vieli et al., 2004), where periods of high melt coincided with reduced seasonal acceleration or even deceleration. Furthermore, numerical modelling results from HH (Nick et al., 2009) suggest that changes in frontal position, as opposed to meltwater-enhanced basal lubrication, are the dominant control on interannual behaviour. Thus, evidence suggests that meltwater-enhanced basal lubrication may significantly influence marine-terminating outlet glacier dynamics at subannual timescales, but its role in driving interannual retreat remains uncertain.

To date, research into the influence of meltwater on marine-terminating outlet glacier dynamics has predominantly focused on enhanced basal lubrication. However, meltwater may also influence dynamics by promoting crevasse propagation at the terminus and/or lateral margins (Figure 3), which together could reduce resistive stresses and promote glacier retreat (Sohn et al., 1998; van der Veen, 1998; Vieli et al., 2007; Andersen et al., 2010b; van der Veen et al., 2011). This partly agrees with model results from JI, which suggest that increased crevasse water levels can partially reproduce observed patterns of

retreat and acceleration, but this may also reflect the choice of calving model (Vieli and Nick, 2011). Numerical modeling studies also suggest that acceleration at Jakobshavn Isbrae (JI), west Greenland, may have resulted from weakening at its lateral margins, potentially due to hydrofracturing and/or meltwater induced warming of the ice (van der Veen et al., 2011). Thus, whilst the role of meltwater-enhanced fracture as a primary trigger of retreat remains equivocal, this mechanism warrants further consideration given the sensitivity of marine-terminating glaciers to changes at the terminus (Nick et al., 2009; Vieli and Nick, 2011).

4 Subglacial drainage systems of large Arctic ice masses

Research into the subglacial hydrology of Arctic ice masses has predominantly focused on land-terminating sections, but recent advances, particularly from the GIS, may provide insight into the comparative insensitivity of marine-terminating outlet glaciers to meltwater-enhanced basal lubrication at interannual timescales. Although the subglacial hydrology of marine-terminating outlet glaciers is comparatively poorly understood and the response of individual glaciers may vary significantly, observations suggest that the seasonal evolution of the subglacial drainage system is very similar to that observed on temperate, polythermal and land-terminating outlet glaciers and sections of the GIS ablation zone: the subglacial drainage system is thought to evolve during the

melt season, causing variation in the sensitivity of ice velocities to meltwater inputs (Figure 5) (e.g. Bartholomew et al., 2010; Howat et al., 2010; Shepherd et al., 2009; Copland et al., 2003; Vieli et al., 2004; Bartholomew et al., 2011; Sole et al., 2011; Dunse et al., 2012). Early in the melt season, the drainage system may be relatively inefficient (Figure 5) (Kamb, 1987; Bingham et al., 2003; Bartholomew et al., 2010; Price et al., 2008). Consequently, meltwater can rapidly increase basal water pressures, causing rapid ice acceleration and surface uplift (Bartholomew et al., 2010; Copland et al., 2003; Bingham et al., 2005). As the melt season progresses, continued inflow of surface meltwater may promote the development of a more efficient, channellized drainage system which operates at lower basal water pressures (Figure 5) (Kamb, 1987; Bingham et al., 2003; Bingham et al., 2006; Shepherd et al., 2009; Palmer et al., 2011; Sole et al., 2011). Thus, the sensitivity of ice velocities to surface melt may decline and only large meltwater inputs may induce substantial velocity change (Figure 5) (Bartholomew et al., 2010; Shepherd et al., 2009; Schoof, 2010; Dunse et al., 2012). The primary implication of these results is that ice velocities depend not only on surface meltwater inputs, but also on the subglacial hydrological system.

The evolution of the subglacial drainage system has important implications for the response of marine-terminating outlet glaciers to interannual variations in

meltwater availability and atmospheric warming (Sundal et al., 2011; Price et al., 2008; Schoof, 2010; van de Wal et al., 2008). As observed at seasonal timescales, continually high meltwater inputs are likely to promote the formation of an efficient basal drainage system, operating at low water pressures (Figure 5). Consequently, increased meltwater input at interannual timescales may not necessarily equate to increased ice velocities, and may even cause deceleration above critical thresholds of water supply (Schoof, 2010; Sundal et al., 2011; Vieli et al., 2004). This is consistent with empirical results from Kangiata Nunata Sermia, south-western Greenland, where meltwater-induced summer speed-up events are thought to contribute little to annual ice velocities, partly because they are offset by the deceleration associated with the formation of an efficient subglacial system (Sole et al., 2011). The key conclusion of these findings is that the evolution of the hydrological system may act as a buffer against accelerated ice loss through meltwater-enhanced basal sliding in response to increased melt and atmospheric warming (Price et al., 2008; Schoof, 2010; Vieli et al., 2004).

IV Oceanic forcing

Whilst atmospheric warming has received substantial scientific attention, oceanic forcing has been recently recognised as a key control on marine-

terminating outlet glacier dynamics. This was partly instigated by results from the GIS (e.g. Moon and Joughin, 2008; Sole et al., 2008; Pritchard et al., 2009), where retreat rates were approximately two orders of magnitude greater on marine-terminating glaciers (10s to 1000s of m a^{-1}) than on their land-terminating counterparts (0.1 to 1 m a^{-1}) (Figure 7). A similar pattern has been observed elsewhere in the Arctic, including Austfonna ice cap (Dowdeswell et al., 2008), Devon Ice Cap (Burgess and Sharp, 2008; Burgess and Sharp, 2004; Dowdeswell et al., 2004; Shepherd et al., 2007) and in Arctic Alaska (Arendt et al., 2006). Furthermore, thinning rates have been greatest on glaciers occupying deep bedrock troughs (Thomas et al., 2009), which may allow warm, sub-surface Atlantic Water (AW) from the continental shelf to access the glacier termini (e.g. Rignot et al., 2010; Straneo et al., 2010; Straneo et al., 2011). Oceanic forcing may be of particular concern in the near-future, as model predictions suggest that ocean temperatures around the GIS may warm by 1.7 to 2°C by 2100 (Yin et al., 2012).

1 Submarine melting at marine-terminating outlet glacier termini

Measurements of submarine melt rates at the termini of marine-terminating glaciers are rare, but estimates suggest that rates range between 0.7 ± 0.2 and 3.9 ± 0.8 m per day in central west Greenland (Rignot et al., 2010) and 4.34 ± 0.94 m per day at JI (Motyka et al., 2011). Substantially higher melt rates of 6.9

to 12.4 m per day have been estimated at LeConte Glacier, Alaska (Figure 1) (Motyka et al., 2003), probably reflecting its comparatively southerly location. These results highlight the potential sensitivity of marine-terminating glaciers to oceanic warming, which could influence outlet glacier dynamics via a number of mechanisms (Figure 8). First, enhanced submarine melting may cause grounding-line retreat at floating and grounded margins, potentially resulting in further un-grounding and the development of positive feedbacks if retreat occurs into deeper water (Vieli et al., 2001; Meier and Post, 1987; Howat et al., 2008a; Joughin et al., 2008b; Vieli and Nick, 2011; Nick et al., 2012). Second, oceanic warming may cause rapid thinning of floating termini (e.g. Motyka et al., 2011; Thomas, 2004; Nick et al., 2012) and the formation of deeply incised basal channels (Rignot and Steffen, 2008), which together make the termini more vulnerable to full thickness fracture and eventual disintegration (Figure 8). Third, submarine melting may influence the terminus geometry and calving rates by undercutting at the grounding line and/or waterline (Figure 8) (Vieli et al., 2002; Benn et al., 2007).

2 Oceanic controls on marine-terminating glacier dynamics

Our understanding of oceanic forcing has been largely developed from observations from the GIS, where warming has immediately preceded the retreat and acceleration of a number of marine-terminating outlet glaciers (e.g.

Hanna et al., 2009; Holland et al., 2008; Murray et al., 2010; Motyka et al., 2011; Bevan et al., 2012a; Rignot et al., 2012). This was first investigated in detail at JI, which was one of the earliest and most significant contributors to recent GIS mass losses (Thomas et al., 2003; Motyka et al., 2011; Rignot and Kanagaratnam, 2006; Joughin et al., 2004; Joughin et al., 2008c; Motyka et al., 2010). Following 50 years of comparative stability (Sohn et al., 1998; Csatho et al., 2008), JI's floating terminus began to retreat in October 1998 (Luckman and Murray, 2005) and subsequent periods of acceleration often coincided with the loss of sections of its tongue (Joughin et al., 2004; Joughin et al., 2008c). Initial retreat was accompanied by rapid thinning, which may have ungrounded the tongue from its underlying pinning points, and caused a substantial reduction in resistive stresses (Joughin et al., 2004; Thomas, 2004; Thomas et al., 2003). This may have initiated feedbacks between retreat, dynamic thinning and acceleration, which led to the disintegration of the ice tongue by spring 2003 (Thomas, 2004; Joughin et al., 2004; Joughin et al., 2008c).

The underlying driver(s) of mass losses at JI remain subject to debate, but evidence suggests that oceanic warming, rather than increased air temperatures, was the primary cause (Motyka et al., 2010; Motyka et al., 2011; Holland et al., 2008; Thomas, 2004). Thinning rates on JI's floating tongue far exceeded estimated surface melt rates and closely followed substantial sub-

surface ocean warming, which is thought to have increased basal melt rates by 25% (Motyka et al., 2011; Holland et al., 2008; Thomas et al., 2003). Estimates suggest that the resultant thinning was sufficient to destabilise the ice tongue and to initiate rapid mass loss (Motyka et al., 2011). Numerical modelling results agree with these findings and suggest that increased submarine melting is capable of triggering the behaviour observed at JI, but that dynamic feedbacks are also required (Vieli and Nick, 2011).

Subsequent to retreat at JI, marine-terminating outlet glaciers in south-eastern Greenland followed a similar progression of dynamic change (e.g. Howat et al., 2008a; Howat et al., 2007; Joughin et al., 2008b; Luckman et al., 2006). Losses began with retreat, thinning and acceleration proportional to retreat, which suggests that changes also resulted from a loss of resistive stresses at the terminus (Howat et al., 2008a; Howat et al., 2007; Howat et al., 2005). The trigger for these changes remains equivocal, with both air temperatures (Hanna et al., 2008; Box et al., 2009) and ocean temperatures (Seale et al., 2011; Hanna et al., 2009; Murray et al., 2010) increasing substantially prior to retreat. However, the initiation of glacier response at the terminus (Howat et al., 2008a; Howat et al., 2005; Howat et al., 2007) suggests that meltwater-enhanced basal lubrication was unlikely to be the primary trigger and that forcing factors operating at the calving front, such as oceanic warming, were the more likely

cause. This is consistent with numerical modelling results from HH, which suggested that interannual glacier dynamics are comparatively insensitive to enhanced basal lubrication, but are acutely sensitive to calving front perturbations (Nick et al., 2009).

3 Marine-terminating outlet glacier dynamics and Atlantic Water distribution

An important emerging theme has been the relationship between marine-terminating outlet glacier dynamics and variations in the distribution and properties of warm Atlantic Water (AW) (Murray et al., 2010; Straneo et al., 2011; Straneo et al., 2010; Holland et al., 2008; Andersen et al., 2012). Until recently, it was assumed that oceanic changes at the continental shelf could be transmitted into outlet glacier fjords, but this was largely untested (Straneo et al., 2010; Mortensen et al., 2011). However, recent studies have shown that AW can access the fjords of a number of large outlet glaciers in Greenland (Straneo et al., 2010; Straneo et al., 2011; Holland et al., 2008; Mayer et al., 2000; Johnson et al., 2011; Christoffersen et al., 2011) and Svalbard (Nilsen et al., 2008). These results marked a significant advance in our understanding, as they demonstrated that rapid connections could exist between marine-terminating outlet glaciers and oceanic variability in the northern North Atlantic, particularly via deep fjords (Straneo et al., 2010). This conclusion was supported by the coincidence of glacier retreat in south-eastern Greenland in

the early 2000s with AW incursion onto the coast (Christoffersen et al., 2011; Murray et al., 2010; Seale et al., 2011) and provides a plausible mechanism for widespread and synchronous retreat.

4 Marine-terminating outlet glacier dynamics and fjord circulation

Recent research into the role of AW has led to increased consideration of the factors controlling its distribution within glacial fjords. A number of possible controls have been identified (Figure 9), including: the temperature, salinity and volume of subtropical waters at the continental shelf; along-shore wind patterns; storm tracks; and fjord stratification (Straneo et al., 2011; Straneo et al., 2010; Nilsen et al., 2008; Christoffersen et al., 2011). Fjord circulation can also be influenced by subglacial meltwater, which forms a rising plume of cool, buoyant water at the calving front and promotes a compensatory inflow of warmer water at depth (Figure 9) (Straneo et al., 2011; Motyka et al., 2011; Motyka et al., 2003). Thus, plumes may substantially increase submarine melt rates (Motyka et al., 2003; Jenkins, 2011; Seale et al., 2011) and model results suggest that melt increases linearly with oceanic warming and to the power of one-third with subglacial discharge (Xu et al., 2012; Jenkins, 2011). A key implication of this relationship is that positive feedbacks could develop, whereby atmospheric warming increases subglacial discharge and ice sheet runoff, which strengthens the plume and enhances submarine melt rates (Seale et al., 2011). Feedbacks

between glacier runoff and ocean properties have been identified as a potential trigger for recent retreat in south-eastern Greenland (Seale et al., 2011; Murray et al., 2010) and variations in meltwater production may be an important control on AW distribution in the region (Murray et al., 2010).

V Sea ice forcing

The increasing focus on oceanic forcing has led to further consideration of the influence of sea ice on marine-terminating Arctic outlet glacier behaviour (Figure 3). Although sea ice is discussed separately, it should be noted that it is influenced by both air and ocean temperatures (Figure 3) and that these factors are not independent. It should also be noted that sea ice concentrations may significantly affect SMB, through their influence on accumulation and ablation patterns (Figure 3) (e.g. Rennermalm et al., 2009; Bamber et al., 2004). The influence of sea ice on marine-terminating Arctic outlet glacier dynamics was first documented in northern Greenland, where semi-permanent fast ice contributed significantly to the stability of several marine-terminating outlet glaciers (Reeh et al., 2001; Mayer et al., 2000; Higgins, 1989; Higgins, 1990; Weidick, 1975). Fast-ice was thought to promote glacier stability by suppressing calving and by preventing calved material from moving away from the terminus (Reeh et al., 2001; Higgins, 1990). In contrast, periods of fast-ice disintegration

were accompanied by rapid calving and release of trapped ice. Early investigations suggested that fast-ice break-up occurred at decadal intervals, when summer temperatures were exceptionally warm (Reeh et al., 2001; Higgins, 1989; Higgins, 1990), but this pattern has changed substantially in recent years, with disintegration now occurring several times per decade (Hughes et al., 2011).

1 Sea ice influence on the seasonal calving cycle

Recent studies have investigated the influence of sea ice on calving rates at more southerly Greenland glaciers (Howat et al., 2010; Ahn and Box, 2010), particularly on JI (Sohn et al., 1998; Joughin et al., 2008c; Amundson et al., 2010). As in northern Greenland, sea ice concentrations at JI appear to influence the timing and nature of calving events, but this occurs on seasonal, as opposed to decadal, timescales (Amundson et al., 2010; Joughin et al., 2008c). In winter, sea ice binds together icebergs to form a semi-rigid, seasonal ice shelf, or *mélange*, which is pushed along the fjord as a coherent mass by the advancing calving front (Figure 10) (Amundson et al., 2010). The *mélange* suppresses calving rates by up to a factor of six and alters the terminus geometry and near-front stress fields, causing seasonal terminus advance and deceleration (Joughin et al., 2008c; Sohn et al., 1998; Amundson et al., 2010). Conversely, spring-time *mélange* disintegration allows high rates of summer

calving to commence, which initiates seasonal retreat and acceleration (Figure 10) (Ahn and Box, 2010; Amundson et al., 2010; Howat et al., 2010; Joughin et al., 2008c). A similar relationship has been documented on the Agassiz Ice Cap, Ellesmere Island, Arctic Canada, where peak glacier velocities have coincided with seasonal sea ice disintegration (Williamson et al., 2008). However, observations also indicated that sea ice weakening and/or thinning, as opposed to complete disintegration, may be sufficient to initiate seasonal acceleration (Williamson et al., 2008).

2 Sea ice influence on interannual marine-terminating outlet glacier behaviour

Observations from JI have contributed substantially to our understanding of sea ice forcing at seasonal timescales, but have also highlighted its potential influence on interannual behaviour of marine-terminating outlet glaciers (Joughin et al., 2008c). Initial retreat at JI began within one year of the onset of sea ice decline in the surrounding Disko Bay (Joughin et al., 2008c). Estimates suggest that the extension of ice free conditions by one or two months may have been sufficient to trigger the initial retreat by extending the duration of seasonally high calving rates (Joughin et al., 2008c). This is consistent with numerical modelling results which demonstrated that reduced mélange duration could trigger rapid retreat at JI, although it could not replicate the magnitude of subsequent seasonal variations in terminus position (Vieli and Nick, 2011). A

similar response has been observed in the Uummannaq region (Howat et al., 2010) and at KG (Christoffersen et al., 2011; Seale et al., 2011), where interannual retreats also followed sea ice decline. It is thought that delayed winter sea ice formation at KG (Christoffersen et al., 2011; Seale et al., 2011) and early mélange clearance in the Uummannaq region (Howat et al., 2010) may have initiated glacier retreat by extending the calving season.

Although the influence of sea ice on marine-terminating outlet glacier behaviour has been little-studied outside of the GIS, Arctic sea ice has declined markedly in recent years (e.g. Rodrigues, 2009; Serreze et al., 2009; Kwok and Rothcock, 2009) and its influence may become increasingly widespread if current losses continue. On the basis of the relationships observed in Greenland, we suggest that sea ice decline may affect glacier dynamics via two potential mechanisms: i), seasonal calving may be extended in areas which currently experience seasonally ice-free conditions; and ii), areas currently characterised by interannual fast-ice may transition to a seasonal sea-ice loss. We suggest that the former process may become increasingly significant on the eastern and central-western Greenland coast, on the western coasts of NZ and Svalbard and in the southern Canadian Arctic, where the ice-free season has extended markedly during the past thirty years (Rodrigues, 2008) and losses are predicted to continue during the 21st century (Figure 11) (ACIA, 2004; IPCC,

2007). This mechanism may eventually cease, however, if areas become perennially ice-free. The latter process may become increasingly important on the coasts of north-eastern Greenland, north-eastern Svalbard, eastern NZ, southern FJL and the northern Canadian Arctic, where sea ice concentrations are predicted to decline markedly by 2100 (Figure 11) (IPCC, 2007; ACIA, 2004). Observations suggest that this may already be occurring in north-eastern Greenland, where fast-ice break up has occurred several times in the past decade (Hughes et al., 2011), in comparison to the decadal intervals recorded by earlier work (Higgins, 1989; Higgins, 1990; Reeh et al., 2001).

VI Key uncertainties and future directions for research

Despite recent advances, the response of marine-terminating outlet glaciers to climatic/oceanic forcing continues to be an area of rapidly developing research and significant uncertainties remain over the relative importance of each forcing factor and the mechanisms by which these factors influence glacier dynamics (Vieli and Nick, 2011; Sole et al., 2008; Howat et al., 2010). The following subsections outline the primary uncertainties surrounding marine-terminating Arctic outlet glacier behaviour and highlight key areas for future research.

1 Spatial variation in the relative importance of climatic/oceanic forcing factors

Our understanding of marine-terminating Arctic outlet glacier response to climatic/oceanic forcing has been primarily based on observations from a small number of Greenland outlet glaciers, with the majority of research focusing on JI and south-eastern Greenland, particularly HH and KG. Consequently, it is uncertain whether the relationships observed at these locations can be extrapolated to other Arctic regions and/or whether recent changes represent a longer-term trend or shorter-term variability (Price et al., 2011; Vieli and Nick, 2011). Although glaciers within certain regions have shown some common response to climatic/oceanic forcing, most notably south-eastern Greenland (Howat et al., 2008a; Murray et al., 2010; Bjørk et al., 2012), this pattern is far from ubiquitous. Results from west Greenland found no correlation between retreat and climatic/oceanic forcing for a sample of 59 marine-terminating outlet glaciers (McFadden et al., 2011) and comparison of 15 major Greenland outlet glaciers between 1985 and 2011 showed some common response to forcing, but also highlighted several notable differences (Bevan et al., 2012a). Furthermore, assessment of decadal and interannual velocity changes on >200 major Greenland outlet glaciers demonstrated substantial variations in glacier behaviour at both regional and local scales and highlighted the importance of glacier-specific factors (Moon et al., 2012). In contrast to the GIS, observations in the Canadian Arctic (Gardner et al., 2011) and Novaya Zemlya (Moholdt et al.,

2012) have found no difference between area-averaged thinning rates in land- and marine-terminating basins (Gardner et al., 2011). Moreover, the longer-term evolution of HH, KG and JI has differed markedly following their earlier mass losses (Howat et al., 2011; Thomas et al., 2011) and numerical modelling studies indicate that marine-terminating outlet glaciers can rapidly adjust to short-term calving front perturbations (Vieli and Nick, 2011). Together, this evidence suggests that the relative importance of climatic/oceanic controls varies across the Arctic and that present theories of outlet glacier response to forcing cannot be universally applied to all glaciers, regions or ice masses. We therefore draw attention to the danger of extrapolating recent rapid mass losses from a small number of glaciers and highlight the need for continued research into the climatic/oceanic drivers of marine-terminating outlet glacier behaviour on each of the major Arctic ice masses.

2 Glacier-specific factors

Results from the GIS have highlighted the substantial variation in marine-terminating outlet glacier response to climatic/oceanic forcing, (McFadden et al., 2011; Moon et al., 2012) and the role of glacier-specific controls, particularly fjord geometry and basal topography, is being increasingly recognised (Howat and Eddy, 2011; Thomas et al., 2009; Joughin et al., 2010; Nick et al., 2009; Joughin et al., 2012; Bevan et al., 2012a). Traditional theories of tidewater

glacier dynamics and ice sheet instability suggest that a reverse basal slope may initiate rapid retreat via a series of positive feedbacks, as the glacier terminus retreats into progressively deeper water (Figure 12) (e.g. Meier and Post, 1987; Vieli et al., 2001; Vieli et al., 2002; Joughin et al., 2008b; Hughes, 1986; Weertman, 1974). This behaviour may occur independently of climatic/oceanic forcing (e.g. Alley, 1991; Pfeffer, 2003), but may also be initiated by perturbations at the calving front (e.g. Meier and Post, 1987; Joughin et al., 2008b; Howat et al., 2008a; Pfeffer, 2007; Nick et al., 2009). However, the influence of overdeepenings on glacier dynamics remains subject to debate and recent modelling results suggest that stable grounding-line positions can be achieved on a reverse bedrock slope (Nick et al., 2010; Gudmundsson et al., 2012). Furthermore, the importance of other glacier-specific factors, such as variations in fjord width, is being increasingly acknowledged (Jamieson et al., in press). Assessing the role of glacier-specific controls is a key area for future study, as inadequate consideration of these factors may lead to substantial errors in estimates of glacier response to climatic/oceanic forcing and their contribution to sea level rise. A full analysis is, however, currently constrained by limited data availability.

3 Quantitative assessment of marine-terminating outlet glacier response to climatic/oceanic forcing

Even on comparatively well-studied sections of the GIS, previous studies have tended to infer causality from the coincidence of climatic/oceanic change and marine-terminating outlet glacier response (e.g. Moon and Joughin, 2008; Luckman et al., 2006). As a consequence, the mechanisms linking climatic/oceanic forcing and glacier dynamics are often poorly understood (Vieli and Nick, 2011; Nick et al., 2009) and the extent to which forcing can explain glacier behaviour has not been extensively assessed. This has been improved in recent years through the development of numerical models focusing on the response of individual outlet glaciers to forcing (Nick et al., 2009; Vieli and Nick, 2011). However, marine-terminating outlet glacier dynamics are not yet adequately represented in ice sheet-scale models (Zwally et al., 2011; Vieli and Nick, 2011; Price et al., 2011) and this is recognised as a significant limitation in our capacity to accurately predict near-future sea level rise (IPCC, 2007). We therefore highlight numerical modelling as an important area for future development and emphasise the need to combine results with remotely sensed and observational data, in order to improve our understanding of recent changes in Arctic marine-terminating outlet glacier dynamics.

VII Conclusions

Arctic ice masses have rapidly lost mass since the mid-1990s due to a combination of negative SMB and accelerated discharge from marine-terminating glaciers (van den Broeke et al., 2009). Studies conducted during the past twenty years have fundamentally altered our understanding of ice mass response to climatic/oceanic forcing and have demonstrated that changes in marine-terminating glacier dynamics can result in dramatic mass losses at annual timescales (e.g. Howat et al., 2008b; Rignot and Kanagaratnam, 2006; Stearns and Hamilton, 2007). In this paper, we identify and review three primary climatic/oceanic drivers of marine-terminating Arctic outlet glacier behaviour: air temperatures, ocean temperatures and sea ice. Although discussed separately, these factors are interconnected and we highlight a number of potentially important linkages which may significantly influence glacier dynamics. We suggest that meltwater-enhanced basal sliding may contribute to marine-terminating outlet glacier velocities at seasonal timescales (Nick et al., 2012; Howat et al., 2010), but its net effect on interannual behaviour may be limited, potentially due to the capacity of the subglacial hydrological system to evolve in response to meltwater inputs (Sundal et al., 2011; Price et al., 2008). Instead, marine-terminating outlet glaciers may respond to atmospheric warming via a number of alternative mechanisms, including: i), hydrofracture of crevasses at the terminus/lateral margins; ii), meltwater-enhanced submarine melting, via

plume circulation and; iii), sea ice loss due to atmospheric warming. Marine-terminating outlet glaciers are potentially highly sensitive to oceanic warming (Rignot et al., 2010), which may cause retreat through: i), submarine melting and rapid thinning across floating sections; ii), grounding-line retreat; iii), alteration of the calving front geometry at the grounding line and/or waterline and; iv), sea ice loss due to oceanic warming. We emphasise the need to further investigate controls on Atlantic Water distribution within glacier fjords and feedbacks between fjord circulation, subglacial meltwater and submarine melting. We also underscore the influence of sea ice on seasonal and interannual outlet glacier dynamics, via its influence on calving rates (Joughin et al., 2008c; Amundson et al., 2010), and suggest that sea ice forcing may become increasingly important during the 21st century if current negative trends continue.

We suggest that the respective role of each climatic/oceanic factor varies across the Arctic and that outlet glacier response to forcing within one region cannot be assumed to apply elsewhere. Moreover, glacier-specific factors may substantially modulate the response of individual glaciers to climatic/oceanic forcing and we highlight this as priority area for future research. Numerical modelling results have improved our understanding of marine-terminating outlet glacier behaviour, but remain a key area for future development. Notwithstanding recent advances, substantial uncertainties remain over the

738 respective roles of the various climatic/oceanic and glacier-specific forcing
739 factors and we highlight the potential danger of extrapolating mass loss rates
740 from a small number of study glaciers. Consequently, the response of marine-
741 terminating Arctic outlet glaciers to climatic/oceanic forcing remains a key area
742 for future research and is crucial for accurate prediction of near-future sea level
743 rise and Arctic ice mass response to climate warming.

744 **Acknowledgements**

745 This work was supported by a Durham Doctoral Studentship granted to J.R.
746 Carr. We thank T. Moon for the provision of Greenland frontal position data and
747 G. Moholdt for helpful comments on Svalbard mass balance data. We thank E.
748 Hanna and N.J. Cox for their assistance with the air temperature data. Two
749 anonymous reviewers are thanked for their constructive comments on the
750 manuscript.

References

- Abdalati W, Krabill W, Frederick E, et al. (2001) Outlet glacier and margin elevation changes: Near-coastal thinning of the Greenland ice sheet. *Journal of Geophysical Research* 106: 33,729–733,741.
- Abdalati W, Krabill W, Frederick E, et al. (2004) Elevation changes of ice caps in the Canadian Arctic Archipelago. *Journal of Geophysical Research* 109: F04007.
- Abdalati W and Steffen K. (2001) Greenland ice sheet melt extent: 1979–1999. *Journal of Geophysical Research* 106: 33983–33988.
- ACIA. (2004) *Impacts of a Warming Arctic: Arctic Climate Impact Assessment*: Cambridge University Press, Cambridge.
- Ahn Y and Box JE. (2010) Glacier velocities from time-lapse photos: technique development and first results from the Extreme Ice Survey (EIS) in Greenland. *Journal of Glaciology* 198: 723–734.
- Alley RB. (1991) Sedimentary processes may cause fluctuations of tidewater glaciers. *Annals of Glaciology* 15: 119–124.
- Alley RB, Dupont TK, Parizek BR, et al. (2005) Access of surface meltwater to beds of sub-freezing glaciers: preliminary insights. *Annals of Glaciology* 40: 8–14.
- Amundson JM, Fahnestock M, Truffer M, et al. (2010) Ice mélange dynamics and implications for terminus stability, Jakobshavn Isbræ, Greenland. *Journal of Geophysical Research* 115: F01005.
- Andersen CS, Larsen TB, Nettles M, et al. (2010a) Spatial and temporal melt variability at Helheim Glacier, East Greenland, and its effect on ice dynamics. *Journal of Geophysical Research* 115: F04041.
- Andersen CS, Straneo F, Hvid M, et al. (2012) Rapid response of Helheim Glacier in Greenland to climate variability over the past century. *Nature Geoscience* 5: 37–41.
- Andersen ML, Larsen TB, Nettles M, et al. (2010b) Spatial and temporal melt variability at Helheim Glacier, East Greenland, and its effect on ice dynamics. *Journal of Geophysical Research* 115: F04041.
- Arendt A, Echelmeyer K, Harrison W, et al. (2006) Updated estimates of glacier volume changes in the western Chugach Mountains, Alaska, and a comparison of regional extrapolation methods. *Journal of Geophysical Research* 111: F03019.
- Arendt A, Luthcke SB, Larsen CF, et al. (2008) Validation of high-resolution GRACE mascon estimates of glacier mass changes in the St Elias Mountains, Alaska, USA, using aircraft laser altimetry. *Journal of Glaciology* 54: 778–787.
- Bamber J, Krabill W, Raper V, et al. (2004) Anomalous recent growth of part of a large Arctic ice cap: Austfonna, Svalbard. *Geophysical Research Letters* 31: L12402.
- Bamber JL, Alley RB and Joughin I. (2007) Rapid response of modern day ice sheets to external forcing. *Earth and Planetary Science Letters* 257: 1–13.
- Bartholomew I, Nienow P, Mair D, et al. (2010) Seasonal evolution of subglacial drainage and acceleration in a Greenland outlet glacier. *Nature Geoscience* 3: 408–411.

- Bartholomew I, Sole A, Mair D, et al. (2011) Supraglacial forcing of subglacial drainage in the ablation zone of the Greenland ice sheet. *Geophysical Research Letters* 38: L08502.
- Bassford RP, Siegert MJ, Dowdeswell J, et al. (2006) Quantifying the Mass Balance of Ice Caps on Severnaya Zemlya, Russian High Arctic. I: Climate and Mass Balance of the Vavilov Ice Cap. *Arctic, Antarctic and Alpine Research* 38: 1-12.
- Benn DI, Warren CR and Mottram RH. (2007) Calving processes and the dynamics of calving glaciers. *Earth Science Reviews* 82: 143-179.
- Bergmann I, Ramillien G and Frappart F. (2012) Climate-driven interannual ice mass evolution in Greenland. *Global and Planetary Change* 82-83: 1-11.
- Bevan SL, Luckman A, Murray T, et al. (2007) Positive mass balance during the late 20th century on Austfonna, Svalbard, revealed using satellite radar interferometry. *Annals of Glaciology* 46: 117-122.
- Bevan SL, Luckman AJ and Murray T. (2012a) Glacier dynamics over the last quarter of a century at Helheim, Kangerdlugssuaq and 14 other major Greenland outlet glaciers. *The Cryosphere* 6: 923-937.
- Bevan SL, Murray T, Luckman A, et al. (2012b) Stable dynamics in a Greenland tidewater glacier over 26 years despite reported thinning. *Annals of Glaciology* 53: 241-248.
- Bhattacharya I, Jezek KC, Wang L, et al. (2009) Surface melt area variability of the Greenland ice sheet: 1979-2008. *Geophysical Research Letters* 36: L20502.
- Bingham RG, Hubbard A, Nienow P, et al. (2008) An investigation into the mechanisms controlling seasonal speedup events at a High Arctic glacier. *Journal of Geophysical Research* 113: F02006.
- Bingham RG, Nienow P and Sharp MJ. (2003) Intra-annual and intra-seasonal flow dynamics of a High Arctic polythermal valley glacier. *Annals of Glaciology* 37: 181-188.
- Bingham RG, Nienow P, Sharp MJ, et al. (2005) Subglacial drainage processes at a High Arctic polythermal valley glacier. *Journal of Glaciology* 51: 15-24.
- Bingham RG, Nienow P, Sharp MJ, et al. (2006) Hydrology and dynamics of a polythermal (mostly cold) High Arctic glacier. *Earth Surface Processes and Landforms* 31: 1463-1479.
- Bjørk AA, Kjær KH, Korsgaard NJ, et al. (2012) An aerial view of 80 years of climate-related glacier fluctuations in southeast Greenland. *Nature Geoscience* Advanced online publication.
- Blaszczyk M, Jania JA and Hagen JM. (2009) Tidewater glaciers of Svalbard: Recent changes and estimates of calving fluxes. *Polish Polar Research* 30: 85-142.
- Bond G, Heinrich H, Broecker W, et al. (1992) Evidence for massive discharges of icebergs into the North Atlantic ocean during the last glacial period. *Nature* 360: 245-249.
- Boon S and Sharp MJ. (2003) The role of hydrologically-driven ice fracture in drainage system evolution on an Arctic glacier. *Geophysical Research Letters* 30: 1916.
- Box JE, Bromwich DH, Veenhuis BA, et al. (2006) Greenland Ice Sheet Surface Mass Balance Variability (1988-2004) from calibrated Polar MM5 Output. *Journal of Climate* 19: 2783-2800.

- Box JE, Yang L, Bromwich DH, et al. (2009) Greenland Ice Sheet Surface Air Temperature Variability: 1840–2007. *Journal of Climate* 22: 4029-4049.
- Briner JP, Bini AC and Anderson RS. (2009) Rapid early Holocene retreat of a Laurentide outlet glacier through an Arctic fjord. *Nature Geoscience* 2: 496-499.
- Burgess DO and Sharp MJ. (2004) Recent Changes in Areal Extent of the Devon Ice Cap, Nunavut, Canada. *Arctic, Antarctic, and Alpine Research* 36: 261-271.
- Burgess DO and Sharp MJ. (2008) Recent changes in thickness of the Devon Island ice cap, Canada. *Journal of Geophysical Research* 113: B07204.
- Cappelen J. (2011) Technical Report 11-05: DMI Monthly Climate Data Collection 1768-2010, Denmark, The Faroe Islands and Greenland In: Cappelen J (ed) *Technical Report 11-05*. Copenhagen: Danish Meteorological Institute.
- Catania GA and Neumann TA. (2010) Persistent englacial drainage features in the Greenland Ice Sheet. *Geophysical Research Letters* 37: L02501.
- Christoffersen P, Mugford R, Heywood KJ, et al. (2011) Warming of waters in an East Greenland fjord prior to glacier retreat: mechanisms and connection to large-scale atmospheric conditions. *The Cryosphere Discussions* 5: 1335-1364.
- Chylek P, Dubey MK and Lesins G. (2006) Greenland warming of 1920–1930 and 1995–2005. *Geophysical Research Letters* 33: L11707.
- Copland L, Sharp MJ and Nienow P. (2003) Links between short-term velocity variations and the subglacial hydrology of a predominantly cold polythermal glacier. *Journal of Glaciology* 49: 337-348.
- Csatho B, Schenk T, van der Veen CJ, et al. (2008) Intermittent thinning of Jakobshavn Isbræ, West Greenland, since the Little Ice Age. *Journal of Glaciology* 54: 131-144.
- Das SB, Joughin I, Behn MD, et al. (2008) Fracture propagation to the base of the Greenland Ice Sheet during supraglacial lake drainage. *Science* 320: 778 – 781.
- Dowdeswell J, Benham TJ, Gorman MR, et al. (2004) Form and flow of the Devon Island Ice Cap, Canadian Arctic. *Journal of Geophysical Research* 109: F02002.
- Dowdeswell J, Benham TJ, Strozzi T, et al. (2008) Iceberg calving flux and mass balance of the Austfonna ice cap on Nordaustlandet, Svalbard. *Journal of Geophysical Research* 113: F03022.
- Dowdeswell J, Hagen JM, Björnsson H, et al. (1997) The Mass Balance of Circum-Arctic Glaciers and Recent Climate Change. *Quaternary Research* 48: 1-14.
- Dunse T, Schuler TV, Hagen JO, et al. (2012) Seasonal speed-up of two outlet glaciers of Austfonna, Svalbard, inferred from continuous GPS measurements. *The Cryosphere* 6: 453-466.
- Dyurgerov MB and McCabe GJ. (2006) Associations between Accelerated Glacier Mass Wastage and Increased Summer Temperature in Coastal Regions. *Arctic, Antarctic and Alpine Research* 38: 190-197.
- Ettema J, Van den Broeke M, van Meijgaard E, et al. (2009) Higher surface mass balance of the Greenland Ice Sheet revealed by high-resolution climate modeling. *Geophysical Research Letters* 36: L12501.

873 Fountain AG and Walder JS. (1998) Water flow through temperate glaciers. *Reviews of*
874 *Geophysics* 36: 299-328.

875 Gardner S, Moholdt G, Wouters B, et al. (2011) Sharply increased mass loss from glaciers and
876 ice caps in the Canadian Arctic Archipelago. *Nature* 473: 357-360.

877 Gerdes R, Larcher MJ, Kauker F, et al. (2003) Causes and development of repeated Arctic
878 Ocean warming events. *Geophysical Research Letters* 30: OCE2.1-OCE2.4

879 Greve R. (2000) On the response of the Greenland Ice Sheet to greenhouse climate change.
880 *Climatic Change* 46: 289-303.

881 Gudmundsson GH, Krug J, Durand G, et al. (2012) The stability of grounding lines on retrograde
882 slopes. *The Cryosphere Discussions* 6: 2597–2619.

883 Hagen JM, Dunse T, Eiken T, et al. (2009) GLACIODYN - The dynamic response of Arctic glaciers
884 to global warming. *American Geophysical Union, Fall Meeting*: Abstract #C53B-06.

885 Hanna E, Cappelen J, Fettweis X, et al. (2009) Hydrologic response of the Greenland Ice Sheet:
886 the role of oceanographic warming. *Hydrological processes* 23: 7-30.

887 Hanna E, Huybrechts P, Steffen K, et al. (2008) Increased runoff from melt from the Greenland
888 Ice Sheet: A response to global warming. *Journal of Climate* 21: 331-341.

889 Higgins AK. (1989) North Greenland ice islands. *Polar Record* 25: 209-212.

890 Higgins AK. (1990) North Greenland glacier velocities and calf ice production. *Polarforschung*
891 60: 1-23.

892 Hodgkins R. (1997) Glacier hydrology in Svalbard, Norwegian High Arctic. *Quaternary Science*
893 *Reviews* 16: 957-973.

894 Holland DM, Thomas RHDY, B., Ribergaard MH, et al. (2008) Acceleration of Jakobshavn Isbræ
895 triggered by warm subsurface ocean waters. *Nature Geoscience* 1: 1-6.

896 Holliday NP, Hughes SL, Bacon S, et al. (2008) Reversal of the 1960s to 1990s freshening trend
897 in the northeast North Atlantic and Nordic Seas. *Geophysical Research Letters* 35:
898 L03614.

899 Howat IM, Ahn Y, Joughin I, et al. (2011) Mass balance of Greenland's three largest outlet
900 glaciers, 2000–2010. *Geophysical Research Letters* 38: L12501.

901 Howat IM, Box JE, Ahn Y, et al. (2010) Seasonal variability in the dynamics of marine-
902 terminating outlet glaciers in Greenland. *Journal of Glaciology* 56: 601-613.

903 Howat IM and Eddy A. (2011) Multi-decadal retreat of Greenland's marine-terminating glaciers.
904 *Journal of Glaciology* 57: 389-396.

905 Howat IM, Joughin I, Fahnestock M, et al. (2008a) Synchronous retreat and acceleration of
906 southeast Greenland outlet glaciers 2000-2006; Ice dynamics and coupling to climate.
907 *Journal of Glaciology* 54: 1-14.

908 Howat IM, Joughin I and Scambos TA. (2007) Rapid changes in ice discharge from Greenland
909 outlet glaciers. *Science* 315: 1559-1561.

910 Howat IM, Joughin I, Tulaczyk S, et al. (2005) Rapid retreat and acceleration of Helheim Glacier,
911 east Greenland. *Geophysical Research Letters* 32: L22502.

912 Howat IM, Smith BE, Joughin I, et al. (2008b) Rates of southeast Greenland ice volume loss
 913 from combined ICESat and ASTER observations. *Geophysical Research Letters* 35:
 914 L17505.
 915 Hughes NE, Wilkinson JP and Wadhams P. (2011) Multi-satellite sensor analysis of fast-ice
 916 development in the Norske Øer Ice Barrier, northeast Greenland. *Annals of Glaciology*
 917 52: 151-160.
 918 Hughes T. (1986) The Jakobshavns effect. *Geophysical Research Letters* 13: 46-48.
 919 Hurrell JW, Kushnir Y, Visbeck MM, et al. (2003) An Overview of the North Atlantic Oscillation.
 920 In: Hurrell JW, Kushnir Y, Ottersen GG, et al. (eds) *The North Atlantic Oscillation:
 921 Climate Significance and Environmental Impact, Geophysical Monograph Series*. 1-35.
 922 Huybrechts P and de Wolde J. (1999) The dynamic response of the Greenland and Antarctic ice
 923 sheets to multiple-century warming. *Journal of Climate* 21: 2169–2188.
 924 Huybrechts P, Letreuilly A and Reeh N. (1991) The Greenland ice sheet and greenhouse
 925 warming. *Palaeogeography, Palaeoclimatology, Palaeoecology* 89: 399-412.
 926 Iken A and Bindenschadler RA. (1986) Combined measurements of subglacial water pressure and
 927 surface velocity of Findelengletscher, Switzerland: Conclusions about drainage system
 928 and sliding mechanism. *Journal of Glaciology* 32: 101-119.
 929 IPCC. (2001) *The Physical Science Basis. Contribution of Working Group I to the Fourth
 930 Assessment Report of the Intergovernmental Panel on Climate Change*: Cambridge
 931 Univ. Press, Cambridge and New York.
 932 IPCC. (2007) *The Physical Science Basis. Contribution of Working Group I to the Fourth
 933 Assessment Report of the Intergovernmental Panel on Climate Change*: Cambridge
 934 Univ. Press, Cambridge and New York.
 935 Jacob T, Wahr J, Pfeffer WT, et al. (2012) Recent contributions of glaciers and ice caps to sea
 936 level rise. *Nature* Advance online publication.
 937 Jamieson SSR, Vieli A, Livingstone SJ, et al. (in press) Ice stream stability on a reverse bed slope.
 938 *Nature Geoscience*.
 939 Jenkins A. (2011) Convection-Driven Melting near the Grounding Lines of Ice Shelves and
 940 Tidewater Glaciers. *Journal of Physical Oceanography* 41: 2279–2294.
 941 Johannessen OM, Khvorostovsky K, Miles MW, et al. (2005) Recent ice-sheet growth in the
 942 interior of Greenland. *Science* 310: 1013 – 1016.
 943 Johnson HL, Münchow A, Falkner KK, et al. (2011) Ocean circulation and properties in
 944 Petermann Fjord, Greenland. *Journal of Geophysical Research* 116: C01003.
 945 Joughin I, Abdalati W and Fahnestock M. (2004) Large fluctuations in speed on Greenland's
 946 Jakobshavn Isbræ glacier. *Nature* 432: 608-610.
 947 Joughin I and Alley RB. (2011) Stability of the West Antarctic ice sheet in a warming world.
 948 *Nature Geoscience* 4: 506-513.
 949 Joughin I, Das SB, King MA, et al. (2008a) Seasonal Speedup Along the Western Flank of the
 950 Greenland Ice Sheet. *Science* 320: 781-783.

951 Joughin I, Howat IM, Alley RB, et al. (2008b) Ice-front variation and tidewater behaviour on
 952 Helheim and Kangerdlugssuaq Glaciers, Greenland. *Journal of Geophysical Research*
 953 113: F01004.
 954 Joughin I, Howat IM, Fahnestock M, et al. (2008c) Continued evolution of Jakobshavn Isbrae
 955 following its rapid speedup. *Journal of Geophysical Research* 113: F04006.
 956 Joughin I, Smith B, Howat IM, et al. (2010) Greenland flow variability from ice-sheet-wide
 957 velocity mapping. *Journal of Glaciology* 56: 415-430.
 958 Joughin I, Smith BE, Howat I, et al. (2012) Seasonal to decadal scale variations in the surface
 959 velocity of Jakobshavn Isbrae, Greenland: Observation and model-based analysis.
 960 *Journal of Geophysical Research* 117: F02030.
 961 Kamb B. (1987) Glacier surge mechanism based on linked cavity configuration of the basal
 962 water conduit system. *Journal of Geophysical Research* 92: 9083-9100.
 963 Khan SA, Wahr J, Bevis M, et al. (2010) Spread of ice mass loss into northwest Greenland
 964 observed by GRACE and GPS. *Geophysical Research Letters* 37: L06501.
 965 Kjær KH, Khan SA, Korsgaard NJ, et al. (2012) Aerial photographs reveal late-20th-Century
 966 dynamic ice loss in northwestern Greenland. *Science* 337: 596-573.
 967 Koerner R. (2005) Mass balance of glaciers in the Queen Elizabeth Islands, Nunavut, Canada.
 968 *Annals of Glaciology* 42: 417-423.
 969 Kotlyakov VM, Glazovskii AF and Frolov IE. (2010) Glaciation in the Arctic. *Herald of the Russian*
 970 *Academy of Sciences* 80: 155-164.
 971 Krabill W, Hanna E, Huybrechts P, et al. (2004) Greenland Ice Sheet: Increased coastal thinning.
 972 *Geophysical Research Letters* 31: L24402.
 973 Krawczynski MJ, Behn MD, Das SB, et al. (2009) Constraints on the lake volume required for
 974 hydro-fracture through ice sheets. *Geophysical Research Letters* 36: L10501.
 975 Kwok R and Rothcock DA. (2009) Decline in Arctic sea ice thickness from submarine and ICESat
 976 records: 1958-2008. *Geophysical Research Letters* 36: L15501.
 977 Luckman A and Murray T. (2005) Seasonal variation in velocity before retreat of Jakobshavn
 978 Isbrae, Greenland. *Geophysical Research Letters* 32: L08501.
 979 Luckman A, Murray T, de Lange R, et al. (2006) Rapid and synchronous ice-dynamic changes in
 980 East Greenland. *Geophysical Research Letters* 33: L03503.
 981 Luthcke SB, Zwally HJ, Abdalati W, et al. (2006) Recent Greenland ice mass loss by drainage
 982 system from satellite gravity observations. *Science* 314: 1286-1289.
 983 Mair D, Burgess DO, Sharp MJ, et al. (2009) Mass balance of the Prince of Wales Icefield,
 984 Ellesmere Island, Nunavut, Canada. *Journal of Geophysical Research* 114: F02011.
 985 Mayer C, Reeh N, Jung-Rothenhäusler F, et al. (2000) The subglacial cavity and implied
 986 dynamics under Nioghalvfjærdsfjorden Glacier, NE-Greenland. *Geophysical Research*
 987 *Letters* 27: 2289-2292.
 988 McFadden EM, Howat IM, Joughin I, et al. (2011) Changes in the dynamics of marine
 989 terminating outlet glaciers in west Greenland (2000-2009). *Journal of Geophysical*
 990 *Research* 116: F02022.

991 Meier MF, Dyurgerov MB, Rick UK, et al. (2007) Glaciers Dominate Eustatic Sea-Level Rise in
 992 the 21st Century. . *Science* 317: 1064-1067.
 993 Meier MF and Post A. (1987) Fast tidewater glaciers. *Journal of Geophysical Research* 92:
 994 9051–9058.
 995 Mémin A, Rogister Y, Hinderer J, et al. (2011) Secular gravity variation at Svalbard (Norway)
 996 from ground observations and GRACE satellite data. *Geophysical Journal International*
 997 184: 1119-1130.
 998 Moholdt G, Hagen JM, Eiken T, et al. (2010a) Geometric changes and mass balance of the
 999 Austfonna ice cap, Svalbard. *The Cryosphere* 4: 21-34.
 1000 Moholdt G, Nuth C, Hagen JO, et al. (2010b) Recent elevation changes of Svalbard glaciers
 1001 derived from ICESat laser altimetry. *Remote Sensing of Environment* 114: 2756–2767.
 1002 Moholdt G, Wouters B and Gardner AS. (2012) Recent mass changes of glaciers in the Russian
 1003 High Arctic. *Geophysical Research Letters* 39: L10502.
 1004 Moon T and Joughin I. (2008) Changes in ice-front position on Greenland’s outlet glaciers from
 1005 1992 to 2007. *Journal of Geophysical Research* 113: F02022.
 1006 Moon T, Joughin I, Smith BE, et al. (2012) 21st-Century evolution of Greenland outlet glacier
 1007 velocities. *Science* 336: 576-578.
 1008 Mortensen J, Lennert K, Bendtsen J, et al. (2011) Heat sources for glacial melt in a sub-Arctic
 1009 fjord (Godthåbsfjord) in contact with the Greenland Ice Sheet. *Journal of Geophysical*
 1010 *Research* 116: C01013.
 1011 Mote TL. (2007) Greenland surface melt trends 1973–2007: Evidence of a large increase in
 1012 2007. *Geophysical Research Letters* 34: L22507.
 1013 Motyka RJ, Fahnestock M and Truffer M. (2010) Volume change of Jakobshavn Isbræ, West
 1014 Greenland: 1985–1997–2007. *Journal of Glaciology* 56: 635-645.
 1015 Motyka RJ, Hunter L, Echelmeyer K, et al. (2003) Submarine melting at the terminus of a
 1016 temperate tidewater glacier, LeConte Glacier, Alaska, U.S.A. *Annals of Glaciology* 36:
 1017 57-65.
 1018 Motyka RJ, Truffer M, Fahnestock M, et al. (2011) Submarine melting of the 1985 Jakobshavn
 1019 Isbræ floating tongue and the triggering of the current retreat. *Journal of Geophysical*
 1020 *Research* 166: F01007.
 1021 Murray T, Scharrer K, James TD, et al. (2010) Ocean regulation hypothesis for glacier dynamics
 1022 in southeast Greenland and implications for ice sheet mass changes. *Journal of*
 1023 *Geophysical Research* 115: F03026.
 1024 Nick FM, Luckman A, Vieli A, et al. (2012) The response of Petermann Glacier, Greenland, to
 1025 large calving events, and its future stability in the context of atmospheric and oceanic
 1026 warming. *Journal of Glaciology* 58: 229 - 239.
 1027 Nick FM, van der Veen CJ, Vieli A, et al. (2010) A physically based calving model applied to
 1028 marine outlet glaciers and implications for the glacier dynamics. *Journal of Geophysical*
 1029 *Research* 56: 781-794.
 1030 Nick FM, Vieli A, Howat IM, et al. (2009) Large-scale changes in Greenland outlet glacier
 1031 dynamics triggered at the terminus. *Nature Geoscience* 2: 110-114.

1032 Nienow P, Sharp MJ and Willis IC. (1998) Seasonal changes in the morphology of the subglacial
1033 drainage system, Haut Glacier d'Arolla, Switzerland. *Earth Surface Processes and*
1034 *Landforms* 23: 825-843.

1035 Nilsen F, Cottier F, Skogseth R, et al. (2008) Fjord– shelf exchanges controlled by ice and brine
1036 production: The interannual variation of Atlantic Water in Isfjorden, Svalbard.
1037 *Continental Shelf Research* 28: 1838– 1853.

1038 Nuth C, Moholdt G, Kohler J, et al. (2010) Svalbard glacier elevation changes and contribution
1039 to sea level rise. *Journal of Geophysical Research* 115: F01008.

1040 Nuttall A-M and Hodgkins R. (2005) Temporal variations in flow velocity at Finsterwalderbreen,
1041 a Svalbard surge-type glacier. *Annals of Glaciology* 42: 71-76.

1042 Palmer S, Shepherd A, Nienow P, et al. (2011) Seasonal speedup of the Greenland Ice Sheet
1043 linked to routing of surface water. *Earth and Planetary Science Letters* 302: 423-428.

1044 Parizek BR and Alley RB. (2004) Implications of increased Greenland surface melt under global-
1045 warming scenarios: Ice-sheet simulations. *Quaternary Science Reviews* 23: 1013– 1027.

1046 Payne AJ, Vieli A, Shepherd AP, et al. (2004) Recent dramatic thinning of largest West Antarctic
1047 ice stream triggered by oceans. *Geophysical Research Letters* 31: L23401.

1048 Pfeffer WT. (2003) Tidewater glaciers move at their own pace. *Nature* 426: 602.

1049 Pfeffer WT. (2007) A simple mechanism for irreversible tidewater glacier retreat. *Journal of*
1050 *Geophysical Research* 112: F03S25.

1051 Price S, Conway H, Waddington ED, et al. (2008) Model investigations of inland migration of
1052 fast-flowing outlet glaciers and ice streams. *Journal of Glaciology* 54: 49-60.

1053 Price S, Payne AJ, Howat IM, et al. (2011) Committed sea-level rise for the next century from
1054 Greenland ice sheet dynamics during the past decade. *Proceedings of the National*
1055 *Academy of Sciences* 108: 8978-8983.

1056 Pritchard HD, Arthern RJ, Vaughan DG, et al. (2009) Extensive dynamic thinning on the margins
1057 of the Greenland and Antarctic ice sheets. *Nature* 461: 971-975.

1058 Radić V and Hock R. (2011) Regionally differentiated contribution of mountain glaciers and ice
1059 caps to future sea-level rise. *Nature Geoscience* 4
1060 91-94.

1061 Raper V, Bamber J and Krabill W. (2005) Interpretation of the anomalous growth of Austfonna,
1062 Svalbard, a large Arctic ice cap. *Annals of Glaciology* 42: 373-379.

1063 Reeh N, Thomsen HH, Higgins AK, et al. (2001) Sea ice and the stability of north and northeast
1064 Greenland floating glaciers. *Annals of Glaciology* 33: 474-480.

1065 Rennermalm AK, Smith LC, Stroeve JC, et al. (2009) Does sea ice influence Greenland ice sheet
1066 surface-melt? *Environmental Research Letters* 4: 024011.

1067 Rignot E, Box JE, Burgess E, et al. (2008) Mass balance of the Greenland ice sheet from 1958 to
1068 2007. *Geophysical Research Letters* 35: L20502.

1069 Rignot E, Fenty I, Menemenlis D, et al. (2012) Spreading of warm ocean waters around
1070 Greenland as a possible cause for glacier acceleration. *Annals of Glaciology* 53: 257-
1071 266.

1072 Rignot E and Kanagaratnam P. (2006) Changes in the velocity structure of the Greenland Ice
1073 Sheet. *Science* 311: 986–990.

1074 Rignot E, Koppes M and Velicogna I. (2010) Rapid submarine melting of the calving faces of
1075 West Greenland glaciers. *Nature Geoscience* 3: 187-191.

1076 Rignot E and Steffen K. (2008) Channelized bottom melting and stability of floating ice shelves.
1077 *Geophysical Research Letters* 35: L02503.

1078 Rignot E, Velicogna I, Van den Broeke M, et al. (2011) Acceleration of the contribution of the
1079 Greenland and Antarctic ice sheets to sea level rise. *Geophysical Research Letters* 38:
1080 L05503.

1081 Rippin DM, Willis IC, Arnold NS, et al. (2005) Spatial and temporal variations in surface velocity
1082 and basal drag across the tongue of the polythermal glacier midre Lovénbreen,
1083 Svalbard. *Journal of Glaciology* 51: 588-600.

1084 Rodrigues J. (2008) The rapid decline of the sea ice in the Russian Arctic. *Cold Regions Science
1085 and Technology* 54: 124-142.

1086 Rodrigues J. (2009) The increase in the length of the ice-free season in the Arctic. *Cold Regions
1087 Science and Technology* 59: 78–101.

1088 Schoof C. (2010) Ice-sheet acceleration driven by melt supply variability. *Nature* 468: 803-806.

1089 Seale A, Christoffersen P, Mugford R, et al. (2011) Ocean forcing of the Greenland Ice Sheet:
1090 Calving fronts and patterns of retreat identified by automatic satellite monitoring of
1091 eastern outlet glaciers. *Journal of Geophysical Research* 116: F03013.

1092 Serreze MC, Barrett AP, Stroeve JC, et al. (2009) The emergence of surface-based Arctic
1093 amplification. *The Cryosphere* 3: 11-19.

1094 Sharov AI. (2005) Studying changes of ice coasts in the European Arctic. *Geo-Marine letters* 25:
1095 153–166.

1096 Sharov AI. (2010) *Satellite Monitoring and Regional Analysis of Glacier Dynamics in the
1097 Barents-Kara Region*, Graz, Austria: JOANNEUM RESEARCH Forschungsgesellschaft
1098 mbH, Institute of Digital Image Processing.

1099 Sharov AI, Schöner W and R. P. (2009) Spatial features of glacier changes in the Barents-Kara
1100 Sector. *EGU General Assembly* 11: EGU2009-3046.

1101 Shepherd A, Du Z, Benham TJ, et al. (2007) Mass balance of Devon Ice Cap, Canadian Arctic.
1102 *Annals of Glaciology* 46: 249-254.

1103 Shepherd A, Hubbard A, Nienow P, et al. (2009) Greenland ice sheet motion coupled with daily
1104 melting in late summer. *Geophysical Research Letters* 36: L01501.

1105 Sneed WA and Hamilton GS. (2007) Evolution of melt pond volume on the surface of the
1106 Greenland Ice Sheet. *Geophysical Research Letters* 34: L03501.

1107 Sohn HG, Jezek KC and van der Veen CJ. (1998) Jakobshavn Glacier, West Greenland: 30 years
1108 of Spacebourne observations. *Geophysical Research Letters* 25: 2699-2702.

1109 Sole A, Payne T, Bamber J, et al. (2008) Testing hypotheses of the cause of peripheral thinning
1110 of the Greenland Ice Sheet: is land-terminating ice thinning at anomalously high rates?
1111 *The Cryosphere Discussions* 2: 673–710.

1112 Sole AJ, Mair DW, Nienow PW, et al. (2011) Seasonal speedup of a Greenland marine-
 1113 terminating outlet glacier forced by surface melt-induced changes in subglacial
 1114 hydrology. *Journal of Geophysical Research* 116: F03014.

1115 Stearns A and Hamilton GS. (2007) Rapid volume loss from two East Greenland outlet glaciers
 1116 quantified using repeat stereo satellite imagery. *Geophysical Research Letters* 34:
 1117 L05503.

1118 Stern H and Heide-Jørgensen MP. (2003) Trends and variability of sea ice in Baffin Bay and
 1119 Davis Strait, 1953–2001. *Polar Research* 22: 11-18.

1120 Straneo F, Curry RG, Sutherland DA, et al. (2011) Impact of fjord dynamics and glacial runoff on
 1121 the circulation near Helheim Glacier. *Nature Geoscience* Advanced online publication
 1122 (10th April 2011).

1123 Straneo F, Hamilton GS, Sutherland DA, et al. (2010) Rapid circulation of warm subtropical
 1124 waters in a major glacial fjord in East Greenland. *Nature Geoscience* 3: 182-186.

1125 Sundal AV, Shepherd A, Nienow P, et al. (2011) Melt-induced speed-up of Greenland ice sheet
 1126 offset by efficient subglacial drainage. *Nature* 469: 521-524.

1127 Thomas R. (1979) The dynamics of marine ice sheets. *Journal of Glaciology* 24: 167-177.

1128 Thomas R, Frederick E, Li J, et al. (2011) Accelerating ice loss from the fastest Greenland and
 1129 Antarctic glaciers. *Geophysical Research Letters* 38: L10502.

1130 Thomas RH. (2004) Force-perturbation analysis of recent thinning and acceleration of
 1131 Jakobshavn Isbrae, Greenland. *Journal of Glaciology* 50: 57-66.

1132 Thomas RH, Abdalati W, Frederick E, et al. (2003) Investigation of surface melting and dynamic
 1133 thinning on Jakobshavn Isbrae, Greenland. *Journal of Glaciology* 49: 231-239.

1134 Thomas RH, Frederick E, Krabill W, et al. (2008) A comparison of Greenland ice-sheet volume
 1135 changes derived from altimetry measurements. *Journal of Glaciology* 54: 203-212.

1136 Thomas RH, Frederick E, Krabill W, et al. (2006) Progressive increase in ice loss from Greenland.
 1137 *Geophysical Research Letters* 33: L10503.

1138 Thomas RH, Frederick E, Krabill W, et al. (2009) Recent changes on Greenland outlet glaciers.
 1139 *Journal of Glaciology* 55: 147-162.

1140 van de Wal RSW, Boot W, van den Broeke MR, et al. (2008) Large and Rapid Melt-Induced
 1141 Velocity Changes in the Ablation Zone of the Greenland Ice Sheet. *Science* 321: 111 -
 1142 113.

1143 van de Wal RSW and Oerlemans J. (1997) Modelling the short-term response of the Greenland
 1144 ice-sheet to global warming. *Climate Dynamics* 13: 733-744.

1145 van den Broeke M, Bamber J, Ettema J, et al. (2009) Partitioning Recent Greenland Mass Loss.
 1146 *Science* 326: 984-986.

1147 van der Veen CJ. (1998) Fracture mechanics approach to penetration of surface crevasses on
 1148 glaciers *Cold Regions Science and Technology* 27: 31-47.

1149 van der Veen CJ. (2007) Fracture propagation as means of rapidly transferring surface
 1150 meltwater to the base of glaciers. *Geophysical Research Letters* 34: L01501.

1151 van der Veen CJ, Plummer JC and Stearns LA. (2011) Controls on the recent speed-up of
 1152 Jakobshavn Isbræ, West Greenland. *Journal of Glaciology* 57: 770–782.

- Velicogna I. (2009) Increasing rates of ice mass loss from the Greenland and Antarctic ice sheets revealed by GRACE. *Geophysical Research Letters* 36: L19503.
- Velicogna I and Wahr J. (2006) Acceleration of Greenland ice mass loss in spring 2004. *Nature* 443: 329- 331.
- Vieli A, Funk M and Blatter H. (2001) Flow dynamics of tidewater glaciers: a numerical modelling approach. *Journal of Glaciology* 47: 595-606.
- Vieli A, Jania JA, Blatter H, et al. (2004) Short-term velocity variations on Hansbreen, a tidewater glacier in Spitsbergen. *Journal of Glaciology* 50: 389-398.
- Vieli A, Jania JA and Lezek K. (2002) The retreat of a tidewater glacier: observations and model calculations on Hansbreen, Spitsbergen. *Journal of Glaciology* 48: 592-600.
- Vieli A and Nick FM. (2011) Understanding and modelling rapid dynamic changes of tidewater outlet glaciers: issues and implications. *Surveys in Geophysics* 32: 437-485.
- Vieli A, Payne AJ, Shepherd A, et al. (2007) Causes of pre-collapse changes of the Larsen B ice shelf: Numerical modelling and assimilation of satellite observations. *Earth and Planetary Science Letters* 259: 297-306.
- Weertman J. (1974) Stability of the junction of an ice sheet and an ice shelf. *Journal of Glaciology* 13: 3-11.
- Weidick A. (1975) A review of Quaternary investigations in Greenland. *Institute of Polar Studies Report* 55: 161.
- Williamson S, Sharp MJ, Dowdeswell J, et al. (2008) Iceberg calving rates from northern Ellesmere Island ice caps, Canadian Arctic, 1999–2003. *Journal of Glaciology* 54: 391-400.
- Willis IC. (1995) Intra-annual variations in glacier motion: a review. *Progress in Physical Geography* 19: 61-106.
- Winsborrow MCM, Andreassen K, Corner GD, et al. (2010) Deglaciation of a marine-based ice sheet: Late Weichselian palaeo-ice dynamics and retreat in the southern Barents Sea reconstructed from onshore and offshore glacial geomorphology. *Quaternary Science Reviews* 29: 424-442.
- Wouters B, Chambers D and Schrama EJO. (2008) GRACE observes small-scale mass loss in Greenland. *Geophysical Research Letters* 35: L20501.
- Xu Y, Rignot E, Menemenlis D, et al. (2012) Numerical experiments on subaqueous melting of Greenland tidewater glaciers in response to ocean warming and enhanced subglacial discharge. *Annals of Glaciology* 53: 229–234.
- Yin J, Overpeck TJ, Griffies SM, et al. (2012) Different magnitudes of projected subsurface ocean warming around Greenland and Antarctica. *Nature Geoscience* 4: 524-528.
- Zwally HJ, Abdalati W, Herring T, et al. (2002) Surface melt-induced acceleration of Greenland ice-sheet flow. *Science* 297: 218–222.
- Zwally HJ, Giovinetto MB, Jun L, et al. (2005) Mass changes of the Greenland and Antarctic ice sheets and shelves and contributions to sea level rise: 1992-2002. *Journal of Glaciology* 51: 509-527.

1193 Zwally HJ, Li J, Brenner AC, et al. (2011) Greenland ice sheet mass balance: distribution of
1194 increased mass loss with climate warming; 2003–07 versus 1992–2002. *Journal of*
1195 *Glaciology* 57: 88-102.

Region	Sub-region	Rate of mass loss (km ³ a ⁻¹)	Measurement period	Measurement method	Source
Greenland	Greenland Ice Sheet	224.76 ± 19*	1992-2009	SMB /D	Rignot et al., 2011
Greenland	Greenland Ice Sheet	203.57± 8.25* [#]	2003-2010	GRACE	Jacob et al., 2012
Canadian Arctic	Ellesmere, Devon, Axel Heiberg and Baffin islands	56.24 ± 6.42*	2004-2009	SMB/D, ICESat laser altimetry and GRACE	Gardner et al., 2011
Canadian Arctic	Ellesmere, Devon, Axel Heiberg islands	34.23 ± 4.56*	2004-2009	SMB/D, ICESat laser altimetry and GRACE	Gardner et al., 2011
Canadian Arctic	Baffin Island	22.0 ± 4.28*	2004-2009	SMB/D, ICESat laser altimetry and GRACE	Gardner et al., 2011
Russian Arctic	Novaya Zemlya	3.67 ± 2	2003 - 2010	GRACE	Jacob et al., 2012
Russian Arctic	Severnaya Zemlya	0.92 ± 2	2003 - 2010	GRACE	Jacob et al., 2012
Russian Arctic	Franz Josef Land	0 ± 2	2003 - 2010	GRACE	Jacob et al., 2012
Svalbard	Spitzbergen	3.59 ± 1.17	2003-2008	ICESat laser altimetry and SPOT HRS 5 stereoscopic images	Moholdt et al., 2010b
Svalbard	Austfonna Ice Cap	1.3 ± 0.5	2002-2008	ICESat laser altimetry, airborne laser altimetry, GNSS surface profiles and RES	Moholdt et al., 2010a
Svalbard	Barentsoya and Edgeoya	0.46 ± 0.30	2003-2008	ICESat laser altimetry and topographic maps	Moholdt et al., 2010b
Svalbard	Vestfonna Ice Cap	0.39 ± 0.20	2003-2008	ICESat laser altimetry and topographic maps	Moholdt et al., 2010b
Svalbard	Kvitoyjokeln ice cap	0.32 ± 0.08	2003-2008	ICESat laser altimetry and topographic maps	Moholdt et al., 2010b

1196 **Table 1.** Recent mass losses from the major glaciated regions and sub-regions of the Arctic. Data are first ordered according to regional mass loss rates
 1197 and then according to mass loss rates from each sub-region. The most recent estimates of total mass loss were used for each region and the latest
 1198 values obtained from GRACE and SMB/D are presented for the GIS. Abbreviations are as follows: (SMB) Surface mass balance, (D) Discharge, (GRACE)

1199 Gravity Recovery and Climate Experiment, (SPOT) Système Pour l'Observation de la Terre, (GNSS) Global Navigation Satellite System and (RES) Radio
1200 Echo Sounding.* Mass loss rates converted from Gt a^{-1} to $\text{km}^3 \text{a}^{-1}$, assuming an ice density of 0.917 kg km^3 (IPCC, 2007). #This value includes peripheral
1201 ice caps and glaciers (Jacob et al., 2012).




Figure 1. Regional overview map showing the location of major ice masses, outlet glaciers and other sites discussed in the text. Major water masses are also labelled. Glacier abbreviations are as follows: Helheim Glacier (HH), Kangerdlugssuaq Glacier (KG), Dagaard Jensen Gletscher (DJ), Kangiata Nunata Sermia (KNS), Jakobshavn Isbrae (JI), Petermann Glacier (PG), Hansbreen (HB), Duvebreen (DB) and John Evans Glacier (JEG). *Inset:* Overview map of Alaska, showing the location of LeConte Glacier (LCG).




Figure 2. Visible satellite imagery of selected marine-terminating Arctic outlet glaciers and Arctic ice masses at 1:1,000,000 scale. Images are ordered by glacier location, from west to east, and show A) Petermann Glacier, north-west Greenland; B) Kangerdlugssuaq Glacier, east Greenland; C) Vestfonna Ice Cap, Svalbard and; D) Northern ice cap, Novaya Zemlya. Outlet glacier and ice mass locations are shown in Figure 1. Major outlet glaciers are labelled according to terminus type (M = marine; L = land) and approximate near-terminus flow direction is marked (dashed lines). Imagery source: Global Land Cover Facility (www.landcover.org).




Figure 4. Linear trend in mean annual air temperatures between 1990 and 2010 for selected Arctic meteorological stations. Symbol colour shows the magnitude of the linear trend in °C per year between 1990 and 2010. Symbol size shows the R^2 value of the relationship: a larger symbol represents a larger R^2 value and therefore a more statistically significant trend. Meteorological stations were selected according to data availability for the study period. Meteorological data sources: Danish Meteorological Institute, Weather and climate data from Greenland 1958-2010; Norwegian Metrological Institute, Eklima climate database; Royal Netherlands Meteorological Institute, Climate Explorer; Scientific Research Institute of Hydrometeorological Information, World Data Center - Baseline Climatological Data Sets; and National Climate Data and Information Archive, Canadian Daily Climate Data.




Figure 5. Idealised seasonal evolution of glacier response to meltwater inputs. The graph illustrates the theoretical response of outlet glacier velocities to meltwater inputs during the melt season. The bottom panels illustrate an idealised plan view of the subglacial hydrological system at different stages of the melt season (bottom panels modified from Fountain and Walder, 1998). Individual glacier response to meltwater forcing may vary significantly from this idealised situation.




Figure 6. Proposed feedback mechanisms between surface meltwater availability, basal sliding and ice sheet geometry for an idealised section of the GIS. Atmospheric warming may increase surface meltwater input to the bed, resulting in enhanced basal sliding and transfer of a greater portion of the outlet glacier to the ablation zone. Further feedbacks may then develop between dynamic thinning, inland migration of basal sliding and ice acceleration. The response of individual sections of the ice sheet may vary significantly from these idealised theoretical responses.




Figure 7. Mean rate of Greenland outlet glacier frontal position change (m a⁻¹) grouped according to terminus type. Data provided by T. Moon, 2011 (Moon and Joughin, 2008). The mean rate of retreat, advance and net frontal position change were calculated for land-terminating and tidewater glacier termini and are shown in the bars above. Values were calculated for three time periods (1992-2000, 2000-2006 and 2006-2007) and maximum rates of retreat / advance are given in brackets above the corresponding bar. Mean values are calculated from a sample of 139 (1992-2000), 169 (2000-2006) and 154 (2006-2007) tidewater glaciers and 10 (1992-2000), 14 (2000-2006) and 13 (2006-2007) land-terminating glaciers. Glaciers terminating in ice shelves were excluded from the analysis, as data were only available from 3 glaciers for 1992-2000 and 2000-2006 and no data were available for 2006-2007.





Figure 8. Illustration of the influence of oceanic warming and submarine melting on outlet glacier dynamics and geometry for A) an initially floating terminus and B) an initially grounded terminus. In A), feedbacks may develop between submarine melting, grounding line retreat, thinning and calving front retreat. In B), changes in terminus geometry may initiate feedbacks between grounding line/terminus retreat, thinning and floatation.




Figure 9. Schematic illustrating the circulation pattern and water properties within a large Arctic outlet glacier fjord. Fjord circulation and water mass depths are based on conditions within Helheim Glacier fjord (Straneo et al., 2011). The primary controls on fjord circulation are thought to be water properties at the continental shelf, winds/storm tracks and glacial meltwater input.




Figure 10. Illustration of the influence of sea ice and mélange formation on Arctic outlet glacier dynamics during A) mélange formation at the end of the calving season and B) mélange disintegration at the start of the calving season. In A) the mélange binds together material within the fjord, thus suppressing calving and promoting seasonal advance. In B) mélange disintegration allows seasonally high calving rates to commence and promotes glacier retreat.




Figure 11. Multi-model mean sea ice concentration (%) for January to March (JFM) and June to September (JAS) in the Arctic for the periods (a) 1980 to 2000 and b) 2080 to 2100 for the SRES A1B scenario. The dashed white line indicates the present-day 15 % average sea ice concentration limit. Modified from IPCC (2007) and Flato et al., 2004. Note the substantial reduction in summer sea ice concentrations predicted across the Arctic by 2100, which may extend seasonally ice free conditions in southerly areas and may result in a transition from multi-year fast ice to seasonal sea-ice disintegration in northern regions.




Figure 12. Illustration of feedbacks between glacier retreat, dynamic thinning and ice acceleration during retreat into progressively deeper water. Initial retreat reduces resistive stresses acting on the outlet glacier, promoting dynamic thinning and terminus floatation, which in turn makes the terminus increasingly vulnerable to fracture and further retreat. Positive feedbacks may also develop between grounding line retreat and submarine melt rates. These feedbacks may occur independently of climatic/oceanic forcing, but may also be triggered by forcing.